

jc604 U.S. PTO



04/13/00

NEW APPLICATION TRANSMITTAL FORM

jc511 U.S. PTO
09/548409
04/13/00

To the Assistant Commissioner for Patents:

This is a Request for filing a non-provisional patent application under 37 CFR 1.53(b) entitled METHODS AND COMPOSITIONS FOR THE TREATMENT OF PANCREATITIS by the following named inventors:

2	Full Name of Inventor	Last Name: Steward	First Name: Lance	Middle Name: E.	
	Residence and Citizenship	City: Irvine	State or Foreign Country: CALIFORNIA	Country Of Citizenship: USA	
	Post Office Address	Post Office Address: 165 San Leon	City: Irvine	State or Country: CA	Zip Code: 92606
1	Full Name of Inventor	Last Name: SACHS	First Name: GEORGE	Middle Name:	
	Residence and Citizenship	City: ENCINO	State or Foreign Country: CALIFORNIA	Country Of Citizenship: USA	
	Post Office Address	Post Office Address: 17986 BORIS DR.	City: ENCINO	State or Country: CA	Zip Code: 91316
2	Full Name of Inventor	Last Name: AOKI	First Name: KEI	Middle Name: ROGER	
	Residence and Citizenship	City: COTO DE CAZA	State or Foreign Country: CALIFORNIA	Country Of Citizenship: USA	
	Post Office Address	Post Office Address: 2 GINGER LILY COURT	City: COTO DE CAZA	State or Country: CA	Zip Code: 92679

(X) The Commissioner is hereby authorized to use Deposit Account Number 01-0885 for the payment of any extension fees incurred during the prosecution of this application.

(X) Enclosed is a specification of 37 pages, claims 4 pages, abstract 1 page, sequence listing 7 pages.

- (X) Enclosed is an executed oath or declaration.
- () Enclosed is an unsigned oath or declaration.
- (X) A self-addressed return postcard is enclosed for verification of receipt.
- (X) The filing fee is calculated below:

FOR	NUMBER FILED	NUMBER EXTRA	RATE	FEE
Basic Fee (Large entity)			\$760	\$690.00
Total Claims	18 minus 20	= 0	× \$18	.00
Independent Claims	2 minus 3	= 0	× \$78	.00
If application contains any multiple dependent claims, then add \$260.00				
TOTAL FILING FEE				\$690.00

- (X) The Commissioner is hereby authorized to charge the filing fee and excess claim fees (including multiple dependent claim fee) as stated above to Deposit Account No. 01-0885. If this amount is incorrect, or for payment of any other fees that may be incurred as a result of this communication please use said Deposit Account. A duplicate copy of this sheet is enclosed for that purpose.
- (X) A copy of an assignment bestowing all interest in this application to Allergan Sales, Inc is enclosed.
- () New drawings are enclosed in __ sheets.
- (X) A Statement Pursuant to 37 CFR 1.821(f) and a labeled diskette containing the computer readable sequence listing is enclosed.
- () A Statement Pursuant to 37 CFR § 1.821(e), stating that the paper copy and the computer readable form are identical is filed herewith.
- (X) A properly labeled computer readable form of the Sequence Listing accompanies this Application.
- (X) The Power of Attorney in this application is to Carlos A. Fisher, Registration Number 36,510.
- (X) The Power of Attorney appears in the combined Declaration and Power of Attorney, filed herewith.

Steward et al

Docket No: 17282CIP(AOC)

PATENT


() A copy of the Request for Extension of Time filed in the prior application is enclosed.

Please address all future communications to:

Carlos A. Fisher
Registration No. 36,510
ALLERGAN, INC.
T2-7H
2525 Dupont Drive
Irvine, CA 92623
Tel: 714-246-4920
Fax: 714-246-4249

Respectfully submitted,

Date: 4/4/00



Carlos A. Fisher
Registration No. 36,510
Attorney of Record

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Steward et al.

Group Art Unit: Not yet assigned

Serial No.: Not yet assigned

I hereby certify that this correspondence is being deposited with the United States Postal Service as Express Mail bearing Label No. EL079261521US in an envelope addressed to: Assistant Commissioner for Patents, Washington, D.C. 20231 on:

Filed: Herewith

For: Methods and Compositions
For the Treatment of Pancreatitis

Date of Deposit: 4/13/00
Printed Name of Person making Deposit: Bonnie Ferguson

Examiner: Not yet assigned

Signature: Bonnie Ferguson
Date of Signature: 4/13/00

CERTIFICATE OF EXPRESS MAILING

Box Patent Application
Assistant Commissioner for Patents
Washington, D.C. 20231

Dear Sir:

Enclosed are a patent application for filing pursuant to 37 CFR 1.53(b). Specifically, accompanying this communication please find:

- (a) Specification in 37 pages, 4 pages claims, 1 page abstract;
- (b) Transmittal sheet in three (3) pages (in duplicate);
- (c) Signed Declaration and Power of Attorney in three (3) pages;
- (d) Information Disclosure Statement and PTO Form 1449;
- (e) Statement Pursuant to 37 CFR 1.821(f) and a labeled diskette containing the computer readable sequence listing;
- (f) Properly labeled computer readable form of the Sequence List;
- (g) Assignment and Assignment Cover Sheet in five (5) pages.

Respectfully submitted,

Date: 4/13/00

Bonnie Ferguson
Bonnie Ferguson

ALLERGAN, INC.- T2-7H
2525 Dupont Drive
Irvine, CA 92612
Tel: 714-246-4920
Fax: 714-246-4249

5

METHODS AND COMPOSITIONS
FOR THE TREATMENT OF PANCREATITIS

This application is a continuation-in-part of
10 application serial no. 09/288,326, filed April 8, 1999.

Field of the Invention

The present invention includes methods and
15 compositions for the treatment of acute pancreatitis.
In a preferred embodiment the invention concerns the use
of agents to reduce or prevent the secretion of
pancreatic digestive enzymes within the pancreas. Such
agents are targeted to pancreatic cells, and serve to
20 prevent the exocytotic fusion of vesicles containing
these enzymes with the plasma membrane. The invention
is also concerned with methods of treating a mammal
suffering from pancreatitis through the administration
of such agents.

25

Background of the Invention

Pancreatitis is a serious medical condition
involving an inflammation of the pancreas. In acute or
30 chronic pancreatitis the inflammation manifests itself
in the release and activation of pancreatic enzymes
within the organ itself, leading to autodigestion. In
many cases of acute pancreatitis, the condition can lead
to death.

35 In normal mammals, the pancreas, a large gland
similar in structure to the salivary gland, is
responsible for the production and secretion of

5 digestive enzymes, which digest ingested food, and
bicarbonate for the neutralization of the acidic chyme
produced in the stomach. The pancreas contains acinar
cells, responsible for enzyme production, and ductal
cells, which secrete large amounts of sodium bicarbonate
10 solution. The combined secretion product is termed
"pancreatic juice"; this liquid flows through the
pancreatic duct past the sphincter of Oddi into the
duodenum. The secretion of pancreatic juice is
stimulated by the presence of chyme in the upper
15 portions of the small intestine, and the precise
composition of pancreatic juice appears to be influenced
by the types of compounds (carbohydrate, lipid, protein,
and/or nucleic acid) in the chyme.

The constituents of pancreatic juice includes
20 proteases (trypsin, chymotrypsin, carboxypolypeptidase),
nucleases (RNase and DNase), pancreatic amylase, and
lipases (pancreatic lipase, cholesterol esterase and
phospholipase). Many of these enzymes, including the
proteases, are initially synthesized by the acinar cells
25 in an inactive form as zymogens: thus trypsin is
synthesized as trypsinogen, chymotrypsin as
chymotrypsinogen, and carboxypolypeptidase as
procarboxypolypeptidase. These enzymes are activated
according to a cascade, wherein, in the first step,
30 trypsin is activated through proteolytic cleavage by the
enzyme enterokinase. Trypsinogen can also be
autoactivated by trypsin; thus one activation has begun,
the activation process can proceed rapidly. Trypsin, in
turn, activates both chymotrypsinogen and
35 procarboxypolypeptidase to form their active protease
counterparts.

5 The enzymes are normally activated only when they
enter the intestinal mucosa in order to prevent
autodigestion of the pancreas. In order to prevent
premature activation, the acinar cells also co-secrete a
trypsin inhibitor that normally prevents activation of
10 the proteolytic enzymes within the secretory cells and
in the ducts of the pancreas. Inhibition of trypsin
activity also prevents activation of the other
proteases.

Pancreatitis can occur when an excess amount of
15 trypsin saturates the supply of trypsin inhibitor.
This, in turn, can be caused by underproduction of
trypsin inhibitor, or the overabundance of trypsin
within the cells or ducts of the pancreas. In the
latter case, pancreatic trauma or blockage of a duct can
20 lead to localized overabundance of trypsin; under acute
conditions large amounts of pancreatic zymogen secretion
can pool in the damaged areas of the pancreas. If even
a small amount of free trypsin is available activation
of all the zymogenic proteases rapidly occurs, and can
25 lead to digestion of the pancreas (acute pancreatitis)
and in particularly severe cases to the patient's death.

Pancreatic secretion is normally regulated by both
hormonal and nervous mechanisms. When the gastric phase
of stomach secretion occurs, parasympathetic nerve
30 impulses are relayed to the pancreas, which initially
results in acetylcholine release, followed by secretion
of enzymes into the pancreatic acini for temporary
storage.

When acid chyme thereafter enters the small
35 intestine, the mucosal cells of the upper intestine
release a hormone called secretin. In humans, secretin

5 is a 27 amino acid (3400 Dalton) polypeptide initially
produced as the inactive form prosecretin, which is then
activated by proteolytic cleavage. Secretin is then
absorbed into the blood. Secretin causes the pancreas
to secrete large quantities of a fluid containing
10 bicarbonate ion. Secretin does not stimulate the acinar
cells, which produce the digestive enzymes. The
bicarbonate fluid serves to neutralize the chyme and to
provide a slightly alkaline optimal environment for the
enzymes.

15 Another peptide hormone, cholecystokinin (CCK) is
released by the mucosal cells in response to the
presence of food in the upper intestine. As described
in further detail below, human CCK is synthesized as a
protoprotein of 115 amino acids. Active CCK forms are
20 quickly taken into the blood through the digestive
tract, and normally stimulate the secretion of enzymes
by the acinar cells. However, stimulation of the CCK
receptor by the CCK analogs cerulein and CCK-octapeptide
(CCK-8) appears to lead to a worsening of morbidity and
25 mortality in mammals in whom pancreatitis is induced.
See Tani et al., *Pancreas* 5:284-290 (1990).

As indicated above, the digestive enzymes are
synthesized as zymogens; proto-enzyme synthesis occurs
in the rough endoplasmic reticulum of the acinar cells.
30 The zymogens are then packaged within vesicles having a
single lipid bilayer membrane. The zymogens are packed
within the vesicles so densely that they appear as
quasi-crystalline structures when observed under light
microscopy and the zymogen granules are electron-dense
35 when observed under the electron microscope. The
vesicles are localized within the cytoplasm of the

5 acinar cells. Secretion of zymogens by the acinar cells occurs through vesicle docking and subsequent fusion with the plasma membrane, resulting in the liberation of the contents into the extracellular milieu.

Nerve cells appear to secrete
10 neurotransmitters and other intercellular signaling factors through a mechanism of membrane fusion that is shared with other cell types, see e.g., Rizo & Sudhof, *Nature Struct. Biol.* 5:839-842 (October 1998), hereby incorporated by reference herein, including the
15 pancreatic acinar cells.

Although the Applicants do not wish to be bound by theory, it is believed that a vesicle first contacts the intracellular surface of the cellular membrane in a reaction called docking. Following the docking step the
20 membrane fuses with and becomes part of the plasma membrane through a series of steps that currently remain relatively uncharacterized, but which clearly involve certain vesicle and membrane-associated proteins, as has been illustrated using neural models.

25 In neurons, neurotransmitters are packaged within synaptic vesicles, formed within the cytoplasm, then transported to the inner plasma membrane where the vesicles dock and fuse with the plasma membrane. Recent studies of nerve cells employing clostridial neurotoxins
30 as probes of membrane fusion have revealed that fusion of synaptic vesicles with the cell membrane in nerve cells depends upon the presence of specific proteins that are associated with either the vesicle or the target membrane. See *id.* These proteins have been
35 termed SNAREs. As discussed in further detail below, a protein alternatively termed synaptobrevin or VAMP

5 (vesicle-associated membrane protein) is a vesicle-associated SNARE (v-SNARE). There are at least two isoforms of synaptobrevin; these two isoforms are differentially expressed in the mammalian central nervous system, and are selectively associated with
10 synaptic vesicles in neurons and secretory organelles in neuroendocrine cells. The target membrane-associated SNAREs (t-SNAREs) include syntaxin and SNAP-25. Following docking, the VAMP protein forms a core complex with syntaxin and SNAP-25; the formation of the core
15 complex appears to be an essential step to membrane fusion. See Rizo & Sudhof, *id.* and Neimmann et al., *Trends in Cell Biol.* 4:179-185 (May 1994), hereby incorporated by referenced herein.

20 Recently evidence has increasingly indicated that the SNARE system first identified in neural cells is a general model for membrane fusion in eukaryotic cells. A yeast exocytotic core complex similar to that of the synaptic vesicles of mammalian neural cells has been characterized, and found to contain three proteins:
25 Sso 1 (syntaxin 1 homolog), SncI (synaptobrevin homolog), and sec9 (SNAP-25 homolog). Rizo & Sudhof, *id.*

These proteins share a high degree of amino acid sequence homology with their mammalian synaptosomal counterparts.

30 All mammalian non-neuronal cells appear to contain cellubrevin, a synaptobrevin analog - this protein is involved in the intracellular transport of vesicles, and is cleaved by TeTx, BoNT/E, BoNT/F, and BoNT/G. Homologs of syntaxin have been identified in
35 yeast (e.g., sso1p and sso2p) and mammalian non-neuronal cells (syn2p, syn3p, syn4p and syn5p). Finally, as

5 indicated above, a yeast SNAP-25 homolog, sec9 has been identified; this protein appears to essential for vesicle fusion with the plasma membrane.

Intoxication of neural cells by clostridial neurotoxins exploits specific characteristics of the
10 SNARE proteins. These neurotoxins, most commonly found expressed in *Clostridium botulinum* and *Clostridium tetanus*, are highly potent and specific poisons of neural cells. These Gram positive bacteria secrete two related but distinct toxins, each comprising two
15 disulfide-linked amino acid chains: a light chain (L) of about 50 KDa and a heavy chain (H) of about 100 KDa, which are wholly responsible for the symptoms of botulism and tetanus, respectively.

The tetanus and botulinum toxins are among the most
20 lethal substances known to man; both toxins function by inhibiting neurotransmitter release in affected neurons.

The tetanus neurotoxin (TeNT) acts mainly in the central nervous system, while botulinum neurotoxin (BoNT) acts at the neuromuscular junction; both toxins
25 inhibit acetylcholine release from the nerve terminal of the affected neuron into the synapse, resulting in paralysis or reduced target organ function.

The tetanus neurotoxin (TeNT) is known to exist in one immunologically distinct type; the botulinum
30 neurotoxins (BoNT) are known to occur in seven different immunologically distinct serotypes, termed BoNT/A through BoNT/G. While all of these latter types are produced by isolates of *C. botulinum*, two other species, *C. baratii* and *C. butyricum* also produce toxins similar
35 to /F and /E, respectively. See e.g., Coffield et al.,

The Site and Mechanism of Action of Botulinum

5 *Neurotoxin in Therapy with Botulinum Toxin* 3-13
 (Jankovic J. & Hallett M. eds. 1994), the disclosure of
 which is incorporated herein by reference.

 Regardless of type, the molecular mechanism of
 intoxication appears to be similar. In the first step
10 of the process, the toxin binds to the presynaptic
 membrane of the target neuron through a specific
 interaction between the heavy chain and a neuronal cell
 surface receptor; the receptor is thought to be
 different for each type of botulinum toxin and for TeNT.

15 The carboxy terminal (C-terminal) half of the heavy
 chain is required for targeting of the toxin to the cell
 surface. The cell surface receptors, while not yet
 conclusively identified, appear to be distinct for each
 neurotoxin serotype.

20 In the second step, the toxin crosses the plasma
 membrane of the poisoned cell. The toxin is first
 engulfed by the cell through receptor-mediated
 endocytosis, and an endosome containing the toxin is
 formed. The toxin (or light chain thereof) then escapes
25 the endosome into the cytoplasm of the cell. This last
 step is thought to be mediated by the amino terminal (N-
 terminal) half of the heavy chain, which triggers a
 conformational change of the toxin in response to a pH
 of about 5.5 or lower. Endosomes are known to possess a
30 proton pump that decreases intra-endosomal pH. The
 conformational shift exposes hydrophobic residues in the
 toxin, which permits the toxin to embed itself in the
 endosomal membrane. The toxin then translocates through
 the endosomal membrane into the cytosol.

35 Either during or after translocation the disulfide
 bond joining the heavy and light chain is reduced, and

5 the light chain is released into the cytoplasm. The entire toxic activity of botulinum and tetanus toxins is contained in the light chain of the holotoxin; the light chain is a zinc (Zn++) endopeptidase which selectively cleaves the SNARE proteins essential for recognition and
10 docking of neurotransmitter-containing vesicles with the cytoplasmic surface of the plasma membrane, and fusion of the vesicles with the plasma membrane. The light chain of TxNT, BoNT/B, BoNT/D, BoNT/F, and BoNT/G cause specific proteolysis of VAMP, an integral protein.
15 During proteolysis, most of the VAMP present at the cytosolic surface of the synaptic vesicle is inactivated as a result of any one of these cleavage events. Each toxin cleaves a different specific peptide bond.

BoNT/A and /E selectively cleave the plasma
20 membrane-associated SNARE protein SNAP-25; this protein is bound to and present on the cytoplasmic surface of the plasma membrane. BoNT/C1 cleaves syntaxin, which exists as an integral protein having most of its mass exposed to the cytosol. Syntaxin interacts with the
25 calcium channels at presynaptic terminal active zones. See Tonello et al., *Tetanus and Botulism Neurotoxins in Intracellular Protein Catabolism* 251-260 (Suzuki K & Bond J. eds. 1996), the disclosure of which is incorporated by reference as part of this specification.
30 Bo/NTC1 also appears to cleave SNAP-25.

Both TeNT and BoNT are specifically taken up by cells present at the neuromuscular junction. BoNT remains within peripheral neurons and, as indicated above, blocks release of the neurotransmitter
35 acetylcholine from these cells.

By contrast TeNT, through its receptor, enters

5 vesicles that move in a retrograde manner along the axon
to the soma, and is discharged into the intersynaptic
space between motor neurons and the inhibitory neurons
of the spinal cord. At this point, TeNT binds receptors
of the inhibitory neurons, is again internalized, and
10 the light chain enters the cytosol to block the release
of the inhibitory neurotransmitters 4-aminobutyric acid
(GABA) and glycine from these cells. Id.

International Patent Publication No. WO 96/33273
relates to derivatives of botulinum toxin designed to
15 prevent neurotransmitter release from sensory afferent
neurons to treat chronic pain. Such derivatives are
targeted to nociceptive neurons using a targeting moiety
that binds to a binding site of the surface of the
neuron.

20 International Patent Publication No. 98/07864
discusses the production of recombinant toxin fragments
that have domains that enable the polypeptide to
translocate into a target cell or which increase the
solubility of the polypeptide, or both.

25

Summary of the Invention

The present invention concerns methods and
30 compositions useful for the treatment of acute
pancreatitis. This condition is largely due to the
defective secretion of zymogen granules by acinar cells,
and by the premature co-mingling of the secreted
zymogens with lysosomal hydrolysates capable of
35 activating trypsin, thereby triggering the protease
activation cascade and resulting in the destruction of

5 pancreatic tissue.

10 In one embodiment of this aspect, the invention is a therapeutic agent comprising a chimeric protein containing an amino acid sequence-specific endopeptidase activity which will specifically cleave at least one synaptic vesicle-associated protein selected from the group consisting of SNAP-25, syntaxin or VAMP, in combination with the translocation activity of the N-terminus of a clostridial neurotoxin heavy chain, wherein the chimeric protein further comprises a
15 recognition domain which will bind a human cholecystokinin (CCK) receptor. Upon binding of the recognition domain of the protein to the CCK receptor, the protein is specifically transported into cells containing CCK receptors (pancreatic acinar cells)
20 through receptor-mediated endocytosis. In a preferred embodiment, the CCK receptor is the CCK A receptor.

Once inside the acinar cell, the chimeric protein functions in a manner similar to that of a clostridial neurotoxin within its target neuron. The toxin moiety
25 is translocated from the endosome into the cytoplasm, where it acts to cleave a SNARE protein identical or homologous to SNAP-25, syntaxin or VAMP. The cleavage of this protein prevents formation of a core complex between the SNARE proteins and thus prevents or reduces
30 the extent of fusion of the vesicle with the target membrane. This, in turn, results in inhibition of zymogen release from the acinar cells and of zymogen activation by lysosomal hydrolases. The autodigestion of pancreatic tissue in acute pancreatitis is therefore
35 reduced or eliminated.

Another embodiment of the present invention

5 concerns a method of treating a patient suffering from acute pancreatitis by administering an effective amount of such a chimeric protein.

Another embodiment of the invention concerns a therapeutic composition that contains the translocation
10 activity of a clostridial neurotoxin heavy chain in combination with a recognition domain able to bind a specific cell type and a therapeutic element having an activity other than the endopeptidase activity of a clostridial neurotoxin light chain. A non-exclusive list
15 of certain such therapeutic elements includes: hormones and hormone-agonists and antagonists, nucleic acids capable of being used as replication, transcription, or translational templates (e.g., for expression of a protein drug having the desired
20 biological activity or for synthesis of a nucleic acid drug as an antisense agent), enzymes, toxins, and the like.

In a preferred embodiment, the specific cell type is a pancreatic cell, most preferably a pancreatic
25 acinar cell.

Another embodiment is drawn to methods for the treatment of acute pancreatitis comprising contacting an acinar cell with an effective amount of a composition comprising a chimeric protein containing an amino acid
30 sequence-specific endopeptidase activity which will specifically cleave at least one synaptic vesicle-associated protein selected from the group consisting of SNAP-25, syntaxin or VAMP, in combination with the translocation activity of the N-terminus of a
35 clostridial neurotoxin heavy chain, wherein the chimeric protein further comprises a recognition domain able to

5 bind to a cell surface protein characteristic of an
human pancreatic acinar cell. Preferably the cell
surface protein is a CCK receptor protein; most
preferably the protein is the human CCK A protein. CCK
receptors (CCK-A receptor and CCK-B receptor) are found
10 mainly in on the surface of pancreatic acinar cells,
although they are also found in some brain cells and, to
a lesser extent on the surface of gastrointestinal
cells.

Any suitable route of administration may be used in
15 this aspect of the invention. Applicants currently
prefer to administer the therapeutic agent in an
intravenous infusion solution; however methods such as
ingestion (particularly when associated with neurotoxin-
associated proteins (NAPs); see Sharma et al., *J. Nat.*
20 *Toxins* 7:239-253(1998), incorporated by reference
herein), direct delivery to the pancreas, injection and
the like may also be used. The agent is substantially
specifically targeted to pancreatic cells; when the
agent contains a CCK receptor-binding domain, the blood-
25 brain barrier prevents the agent from interacting with
brain cells.

In yet another embodiment the invention provides a
composition comprising a drug or other therapeutic agent
having an activity other than that of a clostridial
30 neurotoxin light chain for intracellular delivery, said
agent joined to the translocation domain of a
clostridial neurotoxin heavy chain and a binding element
able to recognize a cell surface receptor of a target
cell. In a preferred embodiment, the target cell is not
35 a neuron. Also, in this embodiment it is preferred that
the drug or other therapeutic agent has an enzymatic,

5 catalytic, or other self-perpetuating mode of activity,
so that the effective dose of drug is greater than the
number of drug molecules delivered within the target
cell. A non-exclusive list of certain such drugs would
include: hormones and hormone-agonists and antagonists,
10 nucleic acids capable being of being used as
replication, transcription, or translational templates
(e.g., for expression of a protein drug having the
desired biological activity or for synthesis of a
nucleic acid drug as an antisense agent), enzymes,
15 toxins (such as diphtheria toxin or ricin), and the
like.

In this embodiment the drug may be cleavably linked
to the remainder of the composition in such a way as to
allow for the release of the drug from the composition
20 within the target cell.

The presently claimed compositions may be provided
to the patient by intravenous administration, may be
administered during surgery, or may be provided
parenterally.

25 WO 95/32738, which shares ownership with the
present application, describes transport proteins for
the therapeutic treatment of neural cells. This
application is incorporated by reference herein as part
of this specification.

30

Detailed Description of the Preferred Embodiments

In a basic and presently preferred form, the
invention comprises a therapeutic polypeptide comprising
35 three features: a binding element, a translocation
element, and a therapeutic element.

5 The binding element is able to bind to a specific
target cell provided that the target cell is not a motor
neuron or a sensory afferent neuron. Preferably, the
binding element comprises an amino acid chain; also an
independently, it is preferably located at or near the
10 C-terminus of a polypeptide chain. By "binding element"
is meant a chemical moiety able to preferentially bind
to a cell surface marker characteristic of the target
cell under physiological conditions. The cell surface
marker may comprise a polypeptide, a polysaccharide, a
15 lipid, a glycoprotein, a lipoprotein, or may have
structural characteristics of more than one of these.
By "preferentially interact" is meant that the
disassociation constant (K_d) of the binding element for
the cell surface marker is at least one order of
20 magnitude less than that of the binding element for any
other cell surface marker. Preferably, the
disassociation constant is at least 2 orders of
magnitude less, even more preferably the disassociation
constant is at least 3 orders of magnitude less than
25 that of the binding element for any other cell surface
marker to which the therapeutic polypeptide is exposed.
Preferably, the organism to be treated is a human.

In one embodiment the cell surface receptor
comprises the histamine receptor, and the binding
30 element comprises an variable region of an antibody
which will specifically bind the histamine receptor.

In an especially preferred embodiment, the cell
surface marker is a cholecystokinin (CCK) receptor.
Cholecystokinin is a bioactive peptide that functions as
35 both a hormone and a neurotransmitter in a wide variety
of physiological settings. Thus, CCK is involved in the

5 regulation of gall bladder contraction, satiety, gastric emptying, and gut motility; additionally it is involved in the regulation of pancreatic exocrine secretion.

There are two types of CCK receptors, CCK A and CCK B; the amino acid sequences of these receptors have been
10 determined from cloned cDNA. Despite the fact that both receptors are G protein-coupled receptors and share approximately 50% homology, there are distinct differences between their physiological activity. The CCK A receptor is expressed in smooth muscle cells of
15 the gall bladder, smooth muscle and neurons within the gastrointestinal tract, and has a much greater affinity (>10² times higher) for CCK than the related peptide hormone gastrin. The CCK B receptor, found in the stomach and throughout the CNS, has roughly equal
20 ability to bind CCK and gastrin.

The varied activities of CCK can be partly attributed to the fact that CCK is synthesized as procholecystokinin, a protoprotein of 115 amino acids, and is then post-translationally cleaved into a number
25 of active fragments all sharing the same C-terminus. The amino acid sequence of human procholecystokinin is shown below; amino acid residues not present in the biologically active cleavage products are in lower case.

All amino acid sequences herein are shown from N-
30 terminus to C-terminus, unless expressly indicated otherwise:

Human procholecystokinin, having the amino acid sequence SEQ ID NO:1:

35

5 mmsgvclcvlmaavlaagaltqpvpdpagsglqraeeaprrqlr VSQRT
DGESRAHLGA LLARYIQQAR KAPSGRMSIV KNLQNLDP SH
RISDRDYM GW MDF grrsaeeeyeps

Biologically active cleavage products of the full
10 length CCK chain include:
CCK-58, having the amino acid sequence SEQ ID NO:2:

VSQRT DGESRAHLGA LLARYIQQAR KAPSGRMSIV KNLQNLDP SH
RISDRDYM GW MDF;

15 CCK-39, having the amino acid sequence SEQ ID NO:
3:

YIQQAR KAPSGRMSIV KNLQNLDP SH RISDRDYM GW MDF;

20 CCK-33, having the amino acid sequence SEQ ID NO:
4:

KAPSGRMSIV KNLQNLDP SH RISDRDYM GW MDF;

25 CCK-12, having the amino acid sequence SEQ ID NO:
5:

ISDRDYM GW MDF;

30 and CCK-8, having the amino acid sequence SEQ ID
NO: 6:
RDYM GW MDF.

35 In each case, the biologically active polypeptides
contain post-translational modifications; in the case of

5 CCK species binding the CCK-A receptor, amidation of the
C-terminal phenylalanine, and sulfatation of the
tyrosine residue located seven residue from the C-
terminus of the biologically active species are required
for hoigh affinity binding ton the receptor. In the
10 case of CCK-B, only the C-terminal amidation is
necessary; sulfation of the tyrosine appears to make
little diffrence in CCK-B binding. These modifications
appear to be necessary for full biological activity,
although both the unmodified C-terminal pentapeptide and
15 tetrapeptide of CCK retains some biological activity.
Kennedy et al., *J. Biol. Chem.* 272: 2920-2926 (1997),
hereby incorporated by reference herein.

In a preferred embodiment, the biologically active
therapeutic polypeptide of the present invention
20 comprises a CCK binding element containing the post-
translational modifications described above. This
polypeptide can be produced by synthetic chemistry or,
preferably, can be produced by a combination of
recombinant and synthetic means using the "expressed
25 protein ligation" (EPL) method. See Cotton & Muir,
Chemistry & Biology 6:R247 (1999), hereby incorporated
by reference herein. In this method the therapeutic
polypeptide is expressed without the C-terminal binding
element as a fusion protein with an "intein" polypeptide
30 sequence positioned at the C-terminus thereof. The
intein comprises a conserved cysteine, serine, or
threonine residue at its amino terminus; the carboxyl
terminus of the intein contains a functional binding
sequence such as chitin binding domain (CBD), poly His
35 (6 or more consecutive histidine residues), or another
amino acid sequence capable of affinity binding. The

5 coding sequence of this recombinantly expressed polypeptide is constructed using standard recombinant DNA methods.

10 Additionally, standard solid phase peptide synthesis methods are employed to construct a synthetic peptide comprising a C-terminal amidated phenylalanine and the desired CCK amino acid sequence. Such methods are described in e.g., Bodansky, M. and Bodansky, A. *The Practice of Peptide Synthesis* (2d ed. Trost B.M., ed. Springer Laboratory 1994), hereby incorporated by
15 reference herein. The synthetic peptide also contains an sulfated tyrosine at the position 7 residues from the carboxyl terminus. This can be done either by incorporation of commercially available Fmoc-Tyr(OSO₃⁻)-OH into the peptide chain at the 7th amino acid position
20 prior to cleavage of the synthetic peptide from the solid support hereby incorporated by reference herein), or by standard peptide synthesis using tyrosine at position 7, followed by a sulfation reaction of the peptide resulting in tyrosine sulfate at the 7 position.
25 See e.g., Koeller, K.M., *J. Am. Chem. Soc.* 122:742-743 (2000). The synthetic peptide is constructed with a cysteine (or serine or threonine) residue at the amino terminus.

30 It will be understood that one can use either hydroxyl-containing amino acids or cysteine as the amino terminal residue of the intein and the synthetic peptide, and either thiopheol, phenol or another nucleophile capable of creating a reactive ester or thioester linkage in accordance with the expressed
35 protein ligation methods described herein. However,

5 thiol-containing amino acid residues and thipheonol or
another sulfur-containing nucleophile are preferred.

Thus, according to one embodiment of the expressed
protein ligation method, the fusion protein is
immobilized following expression by incubation under
10 selective binding conditions with a surface to which the
binding partner of the carboxyl terminal has been joined
(e.g., where the binding moiety is CBP, the surface may
be a resin to which chitin is conjugated). The
immobilized fusion protein is then permitted to react in
15 a transthioesterification reaction with a S- or O-
containing reagent (such as thiophenol or phenol) and
the synthetic modified peptide described above. In this
step, the intein which is joined to the carboxyl
terminus of the therapeutic polypeptide is cleaved at
20 the thioester (or ester) linkage, thus liberating the
protein from the surface to which it was bound. The
intein may be transiently replaced with the thiophenol
group, and the resulting thioester is then itself
attacked by the cysteine (or serine or threonine)
25 residue of the synthetic peptide; this reaction is then
spontaneously followed by a shift of the carbonyl bond
from S (or O) to the N terminal nitrogen of the
synthetic peptide, to form a peptide bond. The
resultant therapeutic polypeptide thus comprises a
30 threapeutic domain, a translocation domain, and a
binding domain comprising a CCK sequence modified to
contain the naturally occuring post-translational
modifications.

As intended herein, the term "extein" refers to a
35 portion of a chimeric polypeptide that borders one or
more intein, and is subsequently ligated to either

5 another extein or a synthetic polypeptide in the EPL
reaction referred to herein.

As intended herein, the term "intein" refers to a
portion of a chimeric polypeptide containing an N-
terminal cysteine, serine, or threonine which is excised
10 from said polypeptide during the EPL reaction referred
to herein.

Of course, the Applicants contemplate that this
method of producing a CCK-containing therapeutic
polypeptide is exemplary only, and that variations and
15 modification of the above-described method will be well
within the ability and knowledge of those of ordinary
skill in the art in light of the present patent
application.

While it will be understood that the applicants do
20 not wish to be bound by theory, the following findings
may assist an understanding the nature of the
interaction between CCK and the CCK receptors, and thus
between the CCK receptor binding element of an
embodiment of the present invention and its CCK receptor
25 target.

In pancreatic acinar cells the CCK A receptor
undergoes internalization to intracellular sites within
minutes after agonist exposure. Pohl et al., *J. Biol.*
Chem. 272: 18179-18184 (1997), hereby incorporated by
30 reference herein. The CCK B receptor has also shown the
same ligand-dependant internalization response in
transfected NIH 3T3 cells. In the CCK B receptor, but
not the CCK A receptor, the endocytotic feature of the
receptor been shown to be profoundly decreased by the
35 deletion of the C terminal 44 amino acids of the

5 receptor chain, corresponding in both receptors to an cytoplasmic portion of the receptor chain.

Recent studies of the interaction between the CCK A receptor and CCK have shown that the primary receptor sequence region containing amino acid residues 38
10 through 42 is involved in the binding of CCK. Residues Trp₃₉ and Gln₄₀ appear to be essential for the binding of a synthetic CCK C-terminal nonapeptide (in which the methionine residues located at residue 3 and 6 from the C-terminus are substituted by norleucine and threonine
15 respectively) to the receptor. Kennedy et al., *supra*. These residues do not appear to be essential for the binding of CCK analogs JMV 180 (corresponding the synthetic C-terminal heptapeptide of CCK in which the phenylalanylamide residue is substituted by a
20 phenylethyl ester and the threonine is substituted with norleucine), and JMV 179 (in which the phenylalanylamide residue and the L-tryptophan residues of the synthetic CCK nonapeptide are substituted by a phenylethyl ester and D-tryptophan, respectively and the threonine is
25 substituted with norleucine). *Id*.

These and similar studies have shed light on the structure of the CCK A receptor active site. Based on receptor binding experiments, a current structural model indicates that CCK residues Trp₃₀ and Met₃₁ (located at
30 positions 4 and 3, respectively, from the C terminus of mature CCK-8) reside in a hydrophobic pocket formed by receptor residues Leu₃₄₈, Pro₃₅₂, Ile₃₅₃ and Ile₃₅₆. CCK residue Asp₃₂ (located at amino acid position 2 measured from the C terminus of CCK-8) seems to be involved in an
35 ionic interaction with receptor residue Lys₁₁₅. CCK Tyr-

5 sulfate₂₇ (the CCK-8 residue 7 amino acids from C terminus) appears involved in an ionic interaction with receptor residue Lys₁₀₅ and a stacking interaction with receptor residue Phe₁₉₈. Ji, et al., 272 *J. Biol. Chem.* 24393-24401 (1997).

10 Such structural models provide detailed guidance to the person of ordinary skill in the art as to the construction of a variety of binding elements able to retain the binding characteristics of biologically active CCK peptides for the CCK-A receptor, for example,
15 as, for example, by site directed mutagenesis of a clostridial neurotoxin heavy chain. Similarly, models deduced using similar methodologies have been proposed for the CCK B receptor, see e.g., Jagerschmidt, A. et al., *Mol. Pharmacol.* 48:783-789 (1995), and can be used
20 as a basis for the construction of binding elements that retain binding characteristics similar to the CCK B receptor.

It will be appreciated that the CCK-B receptor is known to exist on the surface of neurons associated with
25 the certal nervous system. In one alternative embodiment of the present invention the therapeutic polypeptide may be directed (for example, by intrathecal application) to these neurons rather than to the pancreas); in such a case, the binding element may
30 comprise a CCK containing the C terminal amidation only.

Such a binding element may be constructed using the expressed protein ligation (EPL) methods described above. Indeed, EPL methods may be used to introduce and desired or required modifications to the therapeutic
35 element, the translocation element, and/or the binding element of the claimed therapeutic polypeptide.

5 Additionally, the binding element may comprise a variable region of an antibody which will bind the CCK-A or CCK-B receptor.

 Nucleic acids encoding polypeptides containing such a binding element may be constructed using molecular
10 biology methods well known in the art; see e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory Press 2d ed. 1989), and expressed within a suitable host cell. The disclosure of this latter reference is incorporated by reference
15 herein in its entirety.

 The translocation element comprises a portion of a clostridial neurotoxin heavy chain having a translocation activity. By "translocation" is meant the ability to facilitate the transport of a polypeptide
20 through a vesicular membrane, thereby exposing some or all of the polypeptide to the cytoplasm.

 In the various botulinum neurotoxins translocation is thought to involve an allosteric conformational change of the heavy chain caused by a decrease in pH
25 within the endosome.

 This conformational change appears to involve and be mediated by the N terminal half of the heavy chain and to result in the formation of pores in the vesicular membrane; this change permits the movement of the
30 proteolytic light chain from within the endosomal vesicle into the cytoplasm. See e.g., Lacy, et al., *Nature Struct. Biol.* 5:898-902 (October 1998).

 The amino acid sequence of the translocation-mediating portion of the botulinum neurotoxin heavy
35 chain is known to those of skill in the art; additionally, those amino acid residues within this

5 portion that are known to be essential for conferring
the translocation activity are also known.

It would therefore be well within the ability of
one of ordinary skill in the art, for example, to employ
the naturally occurring N-terminal peptide half of the
10 heavy chain of any of the various *Clostridium tetanus* or
Clostridium botulinum neurotoxin subtypes as a
translocation element, or to design an analogous
translocation element by aligning the primary sequences
of the N-terminal halves of the various heavy chains and
15 selecting a consensus primary translocation sequence
based on conserved amino acid, polarity, steric and
hydrophobicity characteristics between the sequences.
The therapeutic element of the present invention may
comprise, without limitation: active or inactive (i.e.,
20 modified) hormone receptors (such as androgen, estrogen,
retinoid, peroxisome proliferator and ecdysone
receptors etc.), and hormone-agonists and antagonists,
nucleic acids capable of being used as
replication, transcription, or translational templates
25 (e.g., for expression of a protein drug having the
desired biological activity or for synthesis of a
nucleic acid drug as an antisense agent), enzymes,
toxins (including apoptosis-inducing agents), and the
like.

30 In a preferred embodiment, the therapeutic element
is a polypeptide comprising a clostridial neurotoxin
light chain or a portion thereof retaining the SNARE-
protein sequence-specific endopeptidase activity of a
clostridial neurotoxin light chain. The amino acid
35 sequences of the light chain of botulinum neurotoxin
(BoNT) subtypes A-G have been determined, as has the

5 amino acid sequence of the light chain of the tetanus
neurotoxin (TeNT). Each chain contains the Zn⁺⁺-binding
motif **His-Glu-x-x-His** (N terminal direction at the left)
characteristic of Zn⁺⁺-dependent endopeptidases (HELIH
in TeNT, BoNT/A /B and /E; HELNH in BoNT/C; and HELTH in
10 BoNT/D).

Recent studies of the BoNT/A light chain have
revealed certain features important for the activity and
specificity of the toxin towards its target substrate,
SNAP-25. Thus, studies by Zhou et al. *Biochemistry*
15 34:15175-15181 (1995) have indicated that when the light
chain amino acid residue His₂₂₇ is substituted with
tyrosine, the resulting polypeptide is unable to cleave
SNAP-25; Kurazono et al., *J. Biol. Chem.* 14721-14729
(1992) performed studies in the presynaptic cholinergic
20 neurons of the buccal ganglia of *Aplysia californica*
using recombinant BoNT/A light chain that indicated that
the removal of 10 N-terminal or 32 C-terminal residues
did not abolish toxicity, but that removal of 10 N-
terminal or 57 C-terminal residues abolished toxicity in
25 this system. Most recently, the crystal structure of
the entire BoNT/A holotoxin has been solved; the active
site is indicated as involving the participation of
His₂₂₂, Glu₂₂₃, His₂₂₆, Glu₂₆₁ and Tyr₃₆₅. Lacy et al., *supra*.
(These residues correspond to His₂₂₃, Glu₂₂₄, His₂₂₇, Glu₂₆₂
30 and Tyr₃₆₆ of the BoNT/A L chain of Kurazono et al.,
supra.) Interestingly, an alignment of BoNT/A through E
and TeNT light chains reveals that every such chain
invariably has these residues in positions analogous to
BoNT/A. Kurazono et al., *supra*.

5 The catalytic domain of BoNT/A is very specific for the C-terminus of SNAP-25 and appears to require a minimum of 16 SNAP-25 amino acids for cleavage to occur.

 The catalytic site resembles a pocket; when the light chain is linked to the heavy chain via the disulfide bond between Cys₄₂₉ and Cys₄₅₃, the translocation domain of the heavy chain appears to block access to the catalytic pocket until the light chain gains entry to the cytosol. When the disulfide bond is reduced, the two polypeptide chains dissociate, and the catalytic pocket is then "opened" and the light chain is fully active.

 As described above, VAMP and syntaxin are cleaved by BoNT/B, D, F, G and TeNT, and BoNT/C₁, respectively, while SNAP-25 is cleaved by BoNT/A and E.

20 The substrate specificities of the various clostridial neurotoxin light chains other than BoNT/A are known. Therefore, the person of ordinary skill in the art could easily determine the toxin residues essential in these subtypes for cleavage and substrate recognition (for example, by site-directed mutagenesis or deletion of various regions of the toxin molecule followed by testing of proteolytic activity and substrate specificity), and could therefore easily design variants of the native neurotoxin light chain that retain the same or similar activity.

 Additionally, construction of the therapeutic agents set forth in this specification would be easily constructed by the person of skill in the art. It is well known that the clostridial neurotoxins have three functional domains analogous to the three elements of the present invention. For example, and without

5 limitation, the BoNT/A neurotoxin light chain is present
in amino acid residues 1-448 of the BoNT/A prototoxin
(i.e., before nicking of the prototoxin to form the
disulfide-linked dichain holotoxin); this amino acid
sequence is provided below as SEQ ID NO: 7. Active site
10 residues are underlined:

BoNT/A light chain (SEQ ID NO:7)

MPFVNKQFNYKDPVNGVDIAYIKIPNAGQMOPVKAFKIHNKI WV
15 IPERDTFTNP EEGDLNPPPEAKQVPVSYDSTYLSTDNEKDNYLKGVT KLFERIYSTD
LGRMLLTSIVRGIPFWGGSTIDTELKVIDTNCINVIQPDGSYRSEELNLVIIGPSADI
IQFECKSFGHEVLNLTRNGYGSTQYIRFSPDFTFGFEESLEVDTNPLL GAGKFATDPA
VTLAHEL I HAGHRLYGIAINPNRVFKVNTNAYYEMSGLEVSFEELRTFGGHDAKFIDS
LQENEFRLYYNKFKDIASTLNKAKSIVGTTASLQYMKNVFKEKYLLSEDTSGKFSVD
20 KLKFDKLYKMLTEIYTEDNFVKFFKVLNRKTYLNFDAVF KINIVPKVNYTIYDGFNL
RNTNLAANFNGQNT EINNMF TKLKNFTGLFEFYKLLCVRGIITSKTKSLDKGYNK;

The heavy chain N-terminal (H_N) translocation
domain is contained in amino acid residues 449-871 of
25 the BoNT/A amino acid sequence, shown below as SEQ ID
NO: 8; a gated ion channel-forming domain probably
essential for the translocation activity of this peptide
is underlined (see Oblatt-Montal et al., *Protein Sci.*
4:1490-1497(1995), hereby incorporated by reference
30 herein.

ALNDLCIKVNNWDLFFSPSEDNFTNDLNKGEEITS D TNIEAAEENISLDLIQQYYLTFNF
DNEPENISIENLSSDIIGQLELMPNIERFPNGKKYELDKYTMFHYLRAQEF EHGKSRI
ALTNSVNEALLNPSRVYTFFSSDYVKKVNKATEAAMFLGWVEQLVYDFTDETSEVSTT
35 DK IADITIIIPYIGPALNIGNMLYKDDFVGALIFSGAVILLEFIPEIAIPVLGTFALV
SYIANKVLT VQTIDNALS KRNEKWDEVYKYIVTNWLAKVNTQIDLIRKKMKEALENQA
EATKAIINYQYNQYTEEEKNNINFNIDDLSSKLNESINKAMININKFLNQCSVSYLMN
SMIPYGVRKLED F DASLKDALLKYIYDNRGTLIGQVDR LKDKVNNTLSTDIPFQLSKY
VDNQRLSTFT EYIK;

40

5 The heavy chain C-terminal neural cell binding domain is contained in amino acid residues 872-1296 (SEQ ID NO: 9) of the BoNT/A prototoxin.

10 NIINTSILNLRYESNHLIDLSRYASKINIGSKVNFDPIDKNQI
QLFNLESSKIEVILKNAIVYNSMYENFSTSFWIRIPKYFNSISLNNEYTIINCMENNS
GWKVSILNYGEIITLQDTQEIKQRVVFYKYSQMINISDYINRWIFVTITNNRLNNSKIY
INGRLIDQKPISNLGNIHASNNIMFKLDGCRDTHRYIWIWIKYFNLFDKELNEKEIKDLY
DNQSNISGILKDFWGDYLDYDKPYMLNLYDPNKYVDVNNVGIRGYMYLKGPRGSVMTT
15 NIYLNSSLYRGTFI IKKYASGNKDNIVRNNDRVYINVVVKNEYRLATNASQAGVEK
ILSALEIPDVGNLSQVVVMKSKNDQGITNKCKMNLQDNNNGNDIGFIGFHQFNNIKLV
ASNWYNRQIERSSRTLGCSEFIPVDDGWERPL

The amino acid sequence of the BoNT/A prototoxin is encoded by nucleotides 358 to 4245 of the neurotoxin
20 cDNA sequence, set forth herein below as SEQ ID NO: 10.

aagcttctaa atttaaatta ttaagtataa atccaaataa acaatatggt
caaaaacttg
atgaggtaat aatttctgta ttagataata tggaaaaata tatagatata
25 tctgaagata
atagattgca actaatagat aacaaaaata acgcaaagaa gatgataatt
agtaatgata
tatttatctc caattgttta accctatctt ataacggtaa atatatatgt
ttatctatga
30 aagatgaaaa ccataattgg atgatatgta ataattgatat gtcaaagtat
ttgtatttat
ggtcatttaa ataattaata atttaattaa ttttaaatat tataagagggt
gttaaatatg
ccatttgtaa ataaacaatt taattataaa gatcctgtaa atgggtgttga
35 tattgcttat
ataaaaattc caaatgcagg acaaatgcaa ccagtaaaag cttttaaaat
tcataataaa
atatgggtta ttccagaaag agatacattt acaaatcctg aagaaggaga
tttaaatcca
40 ccaccagaag caaaacaagt tccagtttca tattatgatt caacatattt
aagtacagat
aatgaaaaag ataattattt aaaggagggt acaaaattat ttgagagaat
ttattcaact
gatcttgtaa gaatgtgtgt aacatcaata gtaaggggaa taccattttg
45 ggggtggaagt
acaatagata cagaattaaa agttattgat actaattgta ttaatgtgat
acaaccagat
ggtagttata gatcagaaga acttaattcta gtaataatag gaccctcagc
tgatattata

Steward et al

Docket No. 17282CIP(AOC)

Patent

5 cagtttgaat gtaaaagctt tggacatgaa gttttgaatc ttacgcgaaa
 tggttatggc
 tctactcaat acattagatt tagcccagat tttacatttg gttttgagga
 gtcacttgaa
 gttgatacaa atcctctttt aggtgcaggc aaatttgcta cagatccagc
 10 agtaacatta
 gcacatgaac ttatacatgc tggacataga ttatatggaa tagcaattaa
 tccaaatagg
 gtttttaaag taaatactaa tgcctattat gaaatgagtg ggtagaagt
 aagctttgag
 15 gaacttagaa catttggggg acatgatgca aagtttatag atagtttaca
 ggaaaacgaa
 tttcgtctat attattataa taagtttaaa gatatagcaa gtacacttaa
 taaagctaaa
 tcaatagtag gtactactgc ttcattacag tatatgaaaa atgtttttta
 20 agagaaatat
 ctcttatctg aagatacatc tggaaaattt tcggtagata aattaaaatt
 tgataagtta
 tacaaaatgt taacagagat ttacacagag gataattttg ttaagttttt
 taaagtactt
 25 aacagaaaaa catatttgaa ttttgataaa gccgtattta agataaatat
 agtacctaag
 gtaaattaca caatatatga tggatttaat ttaagaaata caaatttagc
 agcaaacttt
 aatgggtcaaa atacagaaat taataatatg aattttacta aactaaaaaa
 30 ttttactgga
 ttgtttgaat ttataagtt gctatgtgta agagggataa taacttctaa
 aactaaatca
 ttagataaag gatacaataa ggcattaaat gatttatgta tcaaagttaa
 taattgggac
 35 ttgtttttta gtccttcaga agataatttt actaatgatc taaataaagg
 agaagaaatt
 acatctgata ctaatataga agcagcagaa gaaaatatta gtttagattt
 aatacacaa
 tattatttaa cctttaattt tgataatgaa cctgaaaata tttcaataga
 40 aaatctttca
 agtgacatta taggccaatt agaacttatg cctaatatag aaagatttcc
 taatggaaaa
 aagtatgagt tagataaata tactatgttc cattatcttc gtgctcaaga
 atttgaacat
 45 ggtaaatcta ggattgcttt aacaaattct gttaacgaag cattattaaa
 tcctagtcgt
 gtttatacat ttttttcttc agactatgta aagaaagtta ataaagctac
 ggaggcagct
 atgttttttag gctgggtaga acaattagta tatgatttta ccgatgaaac
 50 tagcgaagta
 agtactacgg ataaaattgc ggatataact ataattattc catatatagg
 acctgcttta
 aatataggta atatgttata taaagatgat tttgtaggtg ctttaaatatt
 ttcaggagct
 55 gttattctgt tagaatttat accagagatt gcaataacctg tattaggtac
 ttttgcactt

Steward et al

Docket No. 17282CIP(AOC)

Patent

5 gtatcatata ttgcgaataa gggttctaacc gttcaaacaa tagataatgc
 ttttaagtaaa
 agaaatgaaa aatgggatga ggtctataaa tatatagtaa caaatgggtt
 agcaaagggtt
 aatacacaga ttgatctaata aagaaaaaaa atgaaagaag ctttagaaaa
 10 tcaagcagaa
 gcaacaaagg ctataataaa ctatcagtat aatcaatata ctgaggaaga
 gaaaaataat attaatTTTA atattgatga ttttaagttcg aaacttaatg
 agtctataaa taaagctatg attaatataa ataaattttt gaatcaatgc
 tctgtttcat atttaatgaa ttctatgatc
 15 ccttatgggtg ttaaacgggt agaagatttt gatgctagtc ttaaagatgc
 attattaaag
 tatatatatg ataataagagg aactttaatt ggtcaagtag atagattaaa
 agataaagtt
 aataatacac ttagtacaga tatacctttt cagctttcca aatacgtaga
 20 taatcaaaga
 ttattatcta catttactga atatattaag aatattatta atacttctat
 attgaattta
 agatatgaaa gtaatcattt aatagactta tctaggtatg catcaaaaat
 aaatattggt
 25 agtaaagtaa attttgatcc aatagataaa aatcaaattc aattatttaa
 ttttagaaagt
 agtaaaattg aggtaatttt aaaaaatgct attgtatata atagtatgta
 tgaaaatttt
 agtactagct tttggataag aattcctaag tattttaaca gtataagtct
 30 aaataatgaa
 tatacaataa taaattgtat ggaaaataat tcaggatgga aagtatcact
 taattatggt
 gaaataatct ggactttaca ggatactcag gaaataaaaac aaagagtagt
 ttttaaatac
 35 agtcaaataa ttaatatatc agattatata aacagatgga tttttgtaac
 tatcactaat
 aatagattaa ataactctaa aatttatata aatggaagat taatagatca
 aaaaccaatt
 tcaaatttag gtaatatcca tgctagtaat aatataatgt ttaaattaga
 40 tgggtgtaga
 gatacacata gatatatattg gataaaatat tttaatcttt ttgataagga
 attaaatgaa
 aaagaaatca aagatttata tgataatcaa tcaaattcag gtatttttaa
 agacttttgg
 45 ggtgattatt tacaatatga taaaccatac tatatgtaa atttatatga
 tccaaataaa
 tatgtcgatg taaataatgt aggtattaga gggttatatgt atcttaaagg
 gcctagagg
 agcgtaatga ctacaaacat ttattttaaat tcaagtttgt atagggggac
 50 aaaatttatt
 ataaaaaat atgcttctgg aaataaagat aatattgtta gaaataatga
 tctgttatat
 attaatgtag tagttaaaaa taaagaatat aggttagcta ctaatgcac
 acaggcaggc
 55 gtagaaaaaa tactaagtgc attagaaata cctgatgtag gaaatctaag
 tcaagtagta

```

5  gtaatgaagt caaaaaatga tcaaggaata acaaataaat gcaaaatgaa
   tttacaagat
   aataatggga atgatatagg ctttatagga tttcatcagt ttaataatat
   agctaaacta
10  gtagcaagta attggtataa tagacaaata gaaagatcta gtaggacttt
   gggttgctca
   tgggaattta ttcctgtaga tgatggatgg ggagaaaggc cactgtaatt
   aatctcaaac
   tacatgagtc tgtcaagaat tttctgtaaa catccataaa aattttaaaa
   ttaatatggt
15  taagaataac tagatatgag tattgtttga actgcccctg tcaagtagac
   aggtaaaaaa
   ataaaaatta agatactatg gtctgatttc gatattctat cggagtcaga
   ccttttaact
   tttcttgat cctttttgta ttgtaaaact ctatgtattc atcaattgca
20  agttccaatt
   agtcaaaatt atgaaacttt ctaagataat acatttctga ttttataatt
   tcccaaaatc
   cttccatagg accattatca atacatctac caactcgaga catactttga
   gttgcgccta
25  tctcattaag tttattcttg aaagatttac ttgtatattg aaaaccgcta
   tcaactgtgaa
   aaagtggact agcatcagga ttggaggtaa ctgctttatc aaaggtttca
   aagacaagga
   cgttgttatt tgattttcca agtacatagg aaataatgct attatcatgc
30  aaatcaagta
   tttcactcaa gtacgccttt gtttcgtctg ttaac

```

Of course, three distinct domains analogous to those described above for BoNT/A exist for all the BoNT subtypes as well as for TeNT neurotoxin; an alignment of the amino acid sequences of these holotoxins will reveal the sequence coordinates for these other neurotoxin species. Additionally, while sequence information is given above for BoNT/A, the amino acid sequences of all BoNT species and tetanus toxin TeNT are known and can easily be obtained from, for example, the NCBI Gen-Bank Web site: www.ncbi.nlm.nih.gov. The Clostrdial neurotoxin nucleotide and amino acid sequences disclosed at this site are expressly incorporated by reference herein.

5 Preferably, the translocation element and the
binding element of the compositions of the present
invention are separated by a spacer moiety that
facilitates the binding element's binding to the desired
cell surface receptor. Such a spacer may comprise, for
10 example, a portion of the BoNT Hc sequence (so long as
the portion does not retain the ability to bind to the
BoNT or TeNT binding site of motor neurons or sensory
afferent neurons), another sequence of amino acids, or a
hydrocarbon moiety. The spacer moiety may also comprise
15 a proline, serine, threonine and/or cysteine-rich amino
acid sequence similar or identical to a human
immunoglobulin hinge region. In a preferred embodiment,
the spacer region comprises the amino acid sequence of
an immunoglobulin $\gamma 1$ hinge region; such a sequence has
20 the sequence (from N terminus to C terminus):

EPKSCDKTHTCPPCP (SEQ ID NO:11)

It will be understood that none of the examples or
embodiments described herein are to be construed as
limiting the scope of the invention, which is defined
25 solely by the claims that conclude this specification.

Example 1:

An agent for the treatment of acute pancreatitis is
30 constructed as follows.

A culture of *Clostridium botulinum* is permitted to
grown to confluence. The cells are then lysed and total
RNA is extracted according to conventional methods and
in the presence of an RNase inhibitor. The RNA
35 preparation is then passed over a oligo(dT) cellulose
column, the polyadenylated messenger RNA is permitted to

5 bind, and the column is washed with 5-10 column volumes
of 20 mM Tris pH 7.6, 0.5 M NaCl, 1 mM EDTA
(ethylenediamine tetraacetic acid), 0.1% (w/v) SDS
(sodium dodecyl sulfate). Polyadenylated RNA is then
eluted with 2-3 column volumes of STE (10 mM Tris (pH
10 7.6), 1 mM EDTA, 0.05% (w/v) SDS). The pooled mRNA is
then precipitated in 2 volumes of ice cold ethanol,
pelleted in a centrifuge at 10,000 x g for 15 minutes,
then redissolved in a small volume of STE.

The BoNT/A mRNA is used as a template for DNA
15 synthesis using Moloney murine leukemia virus reverse
transcriptase (MMLV-RT), then the L chain and then H_N
chain of the neurotoxin is amplified from the cDNA by
the polymerase chain reaction (PCR) using appropriate
oligonucleotide primers whose sequences are designed
20 based on the BoNT/A neurotoxin cDNA sequence of SEQ ID
NO: 9. These procedures are performed using the
standard techniques of molecular biology as detailed in,
for example, Sambrook et al., already incorporated by
reference herein. The primer defining the beginning of
25 the coding region (5' side of the L chain fragment) is
given a StuI site. The PCR primer defining the 3' end of
the H_N-encoding domain has the following features (from
3' to 5'): a 5' region sufficiently complementary to the
3' end of the H_N-encoding domain to anneal thereto under
30 amplification conditions, a nucleotide sequence encoding
the human immunoglobulin hinge region γ_1 (SEQ ID NO:11),
a nucleotide sequence encoding the human CCK-8
octapeptide (SEQ ID NO:6), and a unique restriction
endonuclease cleavage site.

5 The PCR product (termed BoNT/AL-HN- γ -CCK) is purified
by agarose gel electrophoresis, and cloned into a
pBluescript II SK vector. The resulting plasmid is used
to transform competent *E. coli* cells, and a preparation
of the resulting plasmid is made. The BoNT/AL-HN- γ -CCK
10 fragment is excised from the pBluescript vector and
cloned into a mammalian expression vector immediately
downstream of a strong promoter. The resulting vector
is used to transfect a culture of the appropriate host
cell, which is then grown to confluence. Expression of
15 the BoNT/AL-HN- γ -CCK polypeptide is induced, and the cells
are lysed. The polypeptide is first purified by gel
exclusion chromatography, the fractions containing the
recombinant therapeutic agent are pooled, then the
BoNT/AL-HN- γ -CCK polypeptide is further purified using an
20 anti-Ig affinity column wherein the antibody is directed
to the γ_1 hinge region of a human immunoglobulin.

5 Example 2: Method of Treating a Patient Suffering from
 Acute Pancreatitis

 A therapeutically effective amount of the BoNT/A^{LHN}-
 7-CKK agent constructed and purified as set forth in
10 Example 1 is formulated in an acceptable infusion
 solution. Properties of pharmacologically acceptable
 infusion solutions, including proper electrolyte
 balance, are well known in the art. This solution is
 provided intravenously to a patient suffering from acute
15 pancreatitis on a single day over a period of one to two
 hours. Additionally, the patient is fed intravenously
 on a diet low in complex carbohydrates, complex fats and
 proteins.

 At the beginning of treatment, the patient's
20 pancreas shows signs of autodigestion, as measured by
 blood amylase levels. After the treatment regimen,
 autodigestion has ceased, and the patient's pancreas has
 stabilized.

25 Example 3: Alternative Treatment Method

 In this example, a patient suffering from acute
 pancreatitis is treated as in Example 2, with, the
 therapeutic agent given continuously over a period of
30 two weeks. After the treatment regimen, autodigestion
 has ceased, and the patient's pancreas has stabilized.

5 Example 4: Alternative Treatment Method

In this example, a patient suffering from acute pancreatitis is given a single pharmacologically effective amount of the therapeutic agent of Example 1
10 by parenteral administration. Two days after the treatment regimen, autodigestion has ceased and the patient's pancreas has stabilized.

It will be understood that the present invention is
15 not to be limited by the embodiments and examples described herein, and that the invention is defined solely by the claims that conclude this specification.

5

CLAIMS

What is claimed is:

1. A composition for the treatment of acute
pancreatitis in a mammal comprising,
 - a. a first element comprising a binding
element able to specifically bind a pancreatic
cell surface marker under physiological
conditions,
 - b. a second element comprising a
translocation element able to facilitate the
transfer of a polypeptide across a vesicular
membrane, and
 - c. a third element comprising a therapeutic
element able, when present in the cytoplasm of
a pancreatic cell, to inhibit enzymatic
secretion by said pancreatic cell.
1. The composition of claim 1 wherein said pancreatic
cell is an acinar cell and said cell surface marker
is a CCK receptor.
2. The composition of claim 1 wherein said therapeutic
element will cleave a SNARE protein and cleavage of
said SNARE protein inhibits said secretion.
3. The composition of claim 3 wherein said SNARE
protein is selected from the group consisting of
syntaxin, SNAP-25 and VAMP.

35

- 5 4. The composition of claim 2 wherein said therapeutic element will cleave a SNARE protein, wherein cleavage of said SNARE protein inhibits said secretion.
- 10 5. The composition of claim 5 wherein said SNARE protein is selected from the group consisting of syntaxin, SNAP-25 and VAMP.
- 15 6. The composition of claim 5 wherein said CCK receptor is the human CCK A receptor.
- 20 7. The composition of claim 7 wherein the binding element of said thereapeutic polypeptide comprises a human CCK A amino acid sequence modified by the presence of a C-terminal amidated phenylalanine and a sulfated tyrosine at the position 7 residues from the carboxyl terminus.
- 25 8. The composition of claim 8 wherein said CCK A amino acid sequence comprises SEQ ID NO: 6.
9. The composition of claim 9 wherein said CCK A amino acid sequence comprises SEQ ID NO: 5.
- 30 10. The composition of claim 9 wherein said CCK A amino acid sequence comprises SEQ ID NO: 4.
11. The composition of claim 9 wherein said CCK A amino acid sequence comprises SEQ ID NO: 3.

5 12. The composition of claim 9 wherein said CCK A amino acid sequence comprises SEQ ID NO: 2.

10 13. A method for making a therapeutic polypeptide having a binding element selective for a CCK receptor comprising:

a) expressing within a host cell a recombinant chimeric polypeptide comprising an extein comprising a therapeutic element and a translocational element, and an intein located to the carboxyl terminal side of said extein having at its amino terminus an first amino acid selected from the group consisting of cysteine, serine or threonine,

b) contacting said extein with

20 c) a synthetic peptide comprising a CCK amino acid sequence containing modifications comprising the presence of an amidated phenylalanine at a natural C-terminus of said sequence, and further containing at an N-terminus a second amino acid selected from the group consisting of cysteine, serine or threonine,

25 ii) a nucleophilic reagent able to cause cleavage of said intein from the C-terminus of said extein and the subsequent formation of a peptide bond between said extein C-terminus and the N-terminus of said synthetic peptide through the formation of an activated ester or thioester intermediate.

30

35

- 5 14. The method of claim 14 wherein said first and
second amino acids are cysteine.
15. The method of claim 15 wherein said nucleophilic
reagent is selected from the group consisting of
10 phenol or thiphenol.
16. The method of claim 14 wherein said synthetic
polypeptide further comprises a sulfated tyrosine
at the position 7 amino acids from a natural C
15 terminus of said sequence, and said therapeutic
polypeptide preferentially binds a CCK-A receptor.
17. The method of claim 17 wherein said first and
second amino acids are cysteine.
- 20 18. The method of claim 18 wherein said nucleophilic
reagent is selected from the group consisting of
phenol or thiphenol.

5

ABSTRACT

Methods and compositions for the treatment of acute
pancreatitis in a mammal. Particular compositions
comprise a binding element, a translocation element, and
10 a therapeutic element able to prevent accumulation of
digestive enzymes within the pancreas.

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled **METHODS AND COMPOSITIONS FOR THE TREATMENT OF PANCREATITIS** the specification of which

[illegible]

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, §1.56(a). I hereby claim foreign priority benefits under 35 USC § 119(a)-(d) or §365(b) of any foreign application(s) for patent or inventor's certificate, or §365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the Prior Foreign Applications(s).

Number Country Day/Month/Yr filed) [] Priority Not Claimed

I hereby claim the benefit under 35 USC §119 (e) of any United States provisional application(s) listed below.

Application No.	Filing Date
1	1/1/2020
2	1/1/2020
3	1/1/2020
4	1/1/2020
5	1/1/2020
6	1/1/2020
7	1/1/2020
8	1/1/2020
9	1/1/2020
10	1/1/2020
11	1/1/2020
12	1/1/2020
13	1/1/2020
14	1/1/2020
15	1/1/2020
16	1/1/2020
17	1/1/2020
18	1/1/2020
19	1/1/2020
20	1/1/2020
21	1/1/2020
22	1/1/2020
23	1/1/2020
24	1/1/2020
25	1/1/2020
26	1/1/2020
27	1/1/2020
28	1/1/2020
29	1/1/2020
30	1/1/2020
31	1/1/2020
32	1/1/2020
33	1/1/2020
34	1/1/2020
35	1/1/2020
36	1/1/2020
37	1/1/2020
38	1/1/2020
39	1/1/2020
40	1/1/2020
41	1/1/2020
42	1/1/2020
43	1/1/2020
44	1/1/2020
45	1/1/2020
46	1/1/2020
47	1/1/2020
48	1/1/2020
49	1/1/2020
50	1/1/2020
51	1/1/2020
52	1/1/2020
53	1/1/2020
54	1/1/2020
55	1/1/2020
56	1/1/2020
57	1/1/2020
58	1/1/2020
59	1/1/2020
60	1/1/2020
61	1/1/2020
62	1/1/2020
63	1/1/2020
64	1/1/2020
65	1/1/2020
66	1/1/2020
67	1/1/2020
68	1/1/2020
69	1/1/2020
70	1/1/2020
71	1/1/2020
72	1/1/2020
73	1/1/2020
74	1/1/2020
75	1/1/2020
76	1/1/2020
77	1/1/2020
78	1/1/2020
79	1/1/2020
80	1/1/2020
81	1/1/2020
82	1/1/2020
83	1/1/2020
84	1/1/2020
85	1/1/2020
86	1/1/2020
87	1/1/2020
88	1/1/2020
89	1/1/2020
90	1/1/2020
91	1/1/2020
92	1/1/2020
93	1/1/2020
94	1/1/2020
95	1/1/2020
96	1/1/2020
97	1/1/2020
98	1/1/2020
99	1/1/2020
100	1/1/2020

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application(s) in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

<u>09/288,326</u>	<u>4/8/1999</u>
Application No.	Filing Date



I hereby appoint **CARLOS A. FISHER, Registration No. 36,510** (to whom all communications are to be directed), and the below-named persons (of the same address) individually and collectively my attorneys to prosecute this application and to transact all

business in the Patent and Trademark Office connected therewith and with the resulting patent, with full power to appoint associate attorneys:

<u>Name</u>	<u>Registration No.</u>
Robert Baran	25,806
Stephen Donovan	33,433
Martin A. Voet	25,208

of the following correspondence address: **Allergan, Inc., 2525 Dupont Drive, Irvine, CA. 92612**

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under §1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

FULL NAME OF INVENTOR:			
First Name: Lance	Initial E.	Last Name Steward	
RESIDENCE & CITIZENSHIP			
City Irvine	State or Foreign Country California	Country of Citizenship USA	
POST OFFICE ADDRESS			
Post Office Address 165 San Leon	City Irvine	State or Country CA	Zip Code 92606
SIGNATURE OF INVENTOR 		DATE: 4/6/00	
FULL NAME OF INVENTOR:			
First Name: George	Initial	Last Name Sachs	
RESIDENCE & CITIZENSHIP			
City Encino	State or Foreign Country California	Country of Citizenship USA	
POST OFFICE ADDRESS			
Post Office Address 17986 Boris Drive	City Encino	State or Country CA	Zip Code 91316
SIGNATURE OF INVENTOR 		DATE: 4/13/00	

FULL NAME OF INVENTOR:			
First Name: Kei	Initial Roger	Last Name Aoki	
RESIDENCE & CITIZENSHIP			
City Coto de Caza	State or Foreign Country California	Country of Citizenship USA	
POST OFFICE ADDRESS			
Post Office Address 2 Ginger Lily Court	City Cota de Caza	State or Country CA	Zip Code 92679
SIGNATURE OF INVENTOR <i>Kei Roger Aoki</i>		DATE: 4/5/00	

Steward et al.

DOCKET No. 17282 CIP
PATENT

SEQUENCE LISTING

<110> Steward, Lance E.
Aoki, K. Roger
Sachs, George

<120> Compositions ,and Methods for the
Treatment of Pancreatitis

<130> 17282 CIP

<150> 09/288,326

<151> 1999-04-08

<160> 11

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 129

<212> PRT

<213> Homo sapiens

<400> 1

Ser Glu Gln Glu Asn Cys Glu Leu Ile Ser Thr Ile Asn Gly Met Asn
1 5 10 15
Ser Gly Val Cys Leu Cys Val Leu Met Ala Val Leu Ala Ala Gly Ala
20 25 30
Leu Thr Gln Pro Val Pro Pro Ala Asp Pro Ala Gly Ser Gly Leu Gln
35 40 45
Arg Ala Glu Glu Ala Pro Arg Arg Gln Leu Arg Val Ser Gln Arg Thr
50 55 60
Asp Gly Glu Ser Arg Ala His Leu Gly Ala Leu Leu Ala Arg Tyr Ile
65 70 75 80
Gln Gln Ala Arg Lys Ala Pro Ser Gly Arg Met Ser Ile Val Lys Asn
85 90 95
Leu Gln Asn Leu Asp Pro Ser His Arg Ile Ser Asp Arg Asp Tyr Met
100 105 110
Gly Trp Met Asp Phe Gly Arg Arg Ser Ala Glu Glu Tyr Glu Tyr Pro
115 120 125
Ser

<210> 2

<211> 58

<212> PRT

<213> Homo sapiens

<400> 2

Val Ser Gln Arg Thr Asp Gly Glu Ser Arg Ala His Leu Gly Ala Leu
1 5 10 15
Leu Ala Arg Tyr Ile Gln Gln Ala Arg Lys Ala Pro Ser Gly Arg Met
20 25 30
Ser Ile Val Lys Asn Leu Gln Asn Leu Asp Pro Ser His Arg Ile Ser
35 40 45
Asp Arg Asp Tyr Met Gly Trp Met Asp Phe

50

55

<210> 3
 <211> 39
 <212> PRT
 <213> Homo sapiens

<400> 3
 Tyr Ile Gln Gln Ala Arg Lys Ala Pro Ser Gly Arg Met Ser Ile Val
 1 5 10 15
 Lys Asn Leu Gln Asn Leu Asp Pro Ser His Arg Ile Ser Asp Arg Asp
 20 25 30
 Tyr Met Gly Trp Met Asp Phe
 35

<210> 4
 <211> 33
 <212> PRT
 <213> Homo sapiens

<400> 4
 Lys Ala Pro Ser Gly Arg Met Ser Ile Val Lys Asn Leu Gln Asn Leu
 1 5 10 15
 Asp Pro Ser His Arg Ile Ser Asp Arg Asp Tyr Met Gly Trp Met Asp
 20 25 30
 Phe

<210> 5
 <211> 12
 <212> PRT
 <213> Homo sapiens

<400> 5
 Ile Ser Asp Arg Asp Tyr Met Gly Trp Met Asp Phe
 1 5 10

<210> 6
 <211> 9
 <212> PRT
 <213> Homo sapiens

<400> 6
 Arg Asp Tyr Met Gly Trp Met Asp Phe
 1 5

<210> 7
 <211> 448
 <212> PRT
 <213> Clostridium botulinum

<400> 7
 Met Pro Phe Val Asn Lys Gln Phe Asn Tyr Lys Asp Pro Val Asn Gly
 1 5 10 15
 Val Asp Ile Ala Tyr Ile Lys Ile Pro Asn Ala Gly Gln Met Gln Pro
 20 25 30
 Val Lys Ala Phe Lys Ile His Asn Lys Ile Trp Val Ile Pro Glu Arg

[illegible][illegible]

	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99																																																																																																																																																												
0	00000000	00000001	00000010	00000011	00000100	00000101	00000110	00000111	00001000	00001001	00001010	00001011	00001100	00001101	00001110	00001111	00010000	00010001	00010010	00010011	00010100	00010101	00010110	00010111	00011000	00011001	00011010	00011011	00011100	00011101	00011110	00011111	00100000	00100001	00100010	00100011	00100100	00100101	00100110	00100111	00101000	00101001	00101010	00101011	00101100	00101101	00101110	00101111	00110000	00110001	00110010	00110011	00110100	00110101	00110110	00110111	00111000	00111001	00111010	00111011	00111100	00111101	00111110	00111111	01000000	01000001	01000010	01000011	01000100	01000101	01000110	01000111	01001000	01001001	01001010	01001011	01001100	01001101	01001110	01001111	01010000	01010001	01010010	01010011	01010100	01010101	01010110	01010111	01011000	01011001	01011010	01011011	01011100	01011101	01011110	01011111	01100000	01100001	01100010	01100011	01100100	01100101	01100110	01100111	01101000	01101001	01101010	01101011	01101100	01101101	01101110	01101111	01110000	01110001	01110010	01110011	01110100	01110101	01110110	01110111	01111000	01111001	01111010	01111011	01111100	01111101	01111110	01111111	10000000	10000001	10000010	10000011	10000100	10000101	10000110	10000111	10001000	10001001	10001010	10001011	10001100	10001101	10001110	10001111	10010000	10010001	10010010	10010011	10010100	10010101	10010110	10010111	10011000	10011001	10011010	10011011	10011100	10011101	10011110	10011111	10100000	10100001	10100010	10100011	10100100	10100101	10100110	10100111	10101000	10101001	10101010	10101011	10101100	10101101	10101110	10101111	10110000	10110001	10110010	10110011	10110100	10110101	10110110	10110111	10111000	10111001	10111010	10111011	10111100	10111101	10111110	10111111	11000000	11000001	11000010	11000011	11000100	11000101	11000110	11000111	11001000	11001001	11001010	11001011	11001100	11001101	11001110	11001111	11010000	11010001	11010010	11010011	11010100	11010101	11010110	11010111	11011000	11011001	11011010	11011011	11011100	11011101	11011110	11011111	11100000	11100001	11100010	11100011	11100100	11100101	11100110	11100111	11101000	11101001	11101010	11101011	11101100	11101101	11101110	11101111	11110000	11110001	11110010	11110011	11110100	11110101	11110110	11110111	11111000	11111001	11111010	11111011	11111100	11111101	11111110	11111111

[illegible]

<400> 8

Ala	Leu	Asn	Asp	Leu	Cys	Ile	Lys	Val	Asn	Asn	Trp	Asp	Leu	Phe	Phe
1				5					10					15	
Ser	Pro	Ser	Glu	Asp	Asn	Phe	Thr	Asn	Asp	Leu	Asn	Lys	Gly	Glu	Glu
			20					25					30		
Ile	Thr	Ser	Asp	Thr	Asn	Ile	Glu	Ala	Ala	Glu	Glu	Asn	Ile	Ser	Leu
		35					40					45			
Asp	Leu	Ile	Gln	Gln	Tyr	Tyr	Leu	Thr	Phe	Asn	Phe	Asp	Asn	Glu	Pro
	50					55					60				
Glu	Asn	Ile	Ser	Ile	Glu	Asn	Leu	Ser	Ser	Asp	Ile	Ile	Gly	Gln	Leu
65					70					75					80
Glu	Leu	Met	Pro	Asn	Ile	Glu	Arg	Phe	Pro	Asn	Gly	Lys	Lys	Tyr	Glu
				85					90					95	
Leu	Asp	Lys	Tyr	Thr	Met	Phe	His	Tyr	Leu	Arg	Ala	Gln	Glu	Phe	Glu
			100					105					110		
His	Gly	Lys	Ser	Arg	Ile	Ala	Leu	Thr	Asn	Ser	Val	Asn	Glu	Ala	Leu
	115						120					125			
Leu	Asn	Pro	Ser	Arg	Val	Tyr	Thr	Phe	Phe	Ser	Ser	Asp	Tyr	Val	Lys
	130					135						140			
Lys	Val	Asn	Lys	Ala	Thr	Glu	Ala	Ala	Met	Phe	Leu	Gly	Trp	Val	Glu
145					150					155					160
Gln	Leu	Val	Tyr	Asp	Phe	Thr	Asp	Glu	Thr	Ser	Glu	Val	Ser	Thr	Thr
				165					170					175	
Asp	Lys	Ile	Ala	Asp	Ile	Thr	Ile	Ile	Ile	Pro	Tyr	Ile	Gly	Pro	Ala
			180				185						190		
Leu	Asn	Ile	Gly	Asn	Met	Leu	Tyr	Lys	Asp	Asp	Phe	Val	Gly	Ala	Leu
	195					200					205				
Ile	Phe	Ser	Gly	Ala	Val	Ile	Leu	Leu	Glu	Phe	Ile	Pro	Glu	Ile	Ala
	210					215					220				
Ile	Pro	Val	Leu	Gly	Thr	Phe	Ala	Leu	Val	Ser	Tyr	Ile	Ala	Asn	Lys
225					230					235					240
Val	Leu	Thr	Val	Gln	Thr	Ile	Asp	Asn	Ala	Leu	Ser	Lys	Arg	Asn	Glu
				245					250					255	
Lys	Trp	Asp	Glu	Val	Tyr	Lys	Tyr	Ile	Val	Thr	Asn	Trp	Leu	Ala	Lys
			260					265					270		
Val	Asn	Thr	Gln	Ile	Asp	Leu	Ile	Arg	Lys	Lys	Met	Lys	Glu	Ala	Leu
	275					280						285			
Glu	Asn	Gln	Ala	Glu	Ala	Thr	Lys	Ala	Ile	Ile	Asn	Tyr	Gln	Tyr	Asn
	290					295					300				
Gln	Tyr	Thr	Glu	Glu	Glu	Lys	Asn	Asn	Ile	Asn	Phe	Asn	Ile	Asp	Asp
305					310					315					320
Leu	Ser	Ser	Lys	Leu	Asn	Glu	Ser	Ile	Asn	Lys	Ala	Met	Ile	Asn	Ile
				325					330					335	
Asn	Lys	Phe	Leu	Asn	Gln	Cys	Ser	Val	Ser	Tyr	Leu	Met	Asn	Ser	Met
			340					345					350		
Ile	Pro	Tyr	Gly	Val	Lys	Arg	Leu	Glu	Asp	Phe	Asp	Ala	Ser	Leu	Lys
	355					360						365			
Asp	Ala	Leu	Leu	Lys	Tyr	Ile	Tyr	Asp	Asn	Arg	Gly	Thr	Leu	Ile	Gly
	370					375					380				
Gln	Val	Asp	Arg	Leu	Lys	Asp	Lys	Val	Asn	Asn	Thr	Leu	Ser	Thr	Asp
385					390					395					400
Ile	Pro	Phe	Gln	Leu	Ser	Lys	Tyr	Val	Asp	Asn	Gln	Arg	Leu	Leu	Ser
				405					410					415	
Thr	Phe	Thr	Glu	Tyr	Ile	Lys									
			420												

<210> 9

[illegible][illegible][illegible][illegible][illegible]

<213> Clostridium botulinum

<400> 10

aagcttctaa	atttaaatta	ttaagtataa	atccaaataa	acaatatggt	caaaaacttg	60
atgaggtaat	aattttctgta	ttagataata	tggaaaaata	tatagatata	tctgaagata	120
atagattgca	actaatagat	aacaaaaata	acgcaaagaa	gatgataatt	agtaatgata	180
tattttatttc	caattgttta	accctatctt	ataacggtaa	atatatatgt	ttatctatga	240
aagatgaaaa	ccataattgg	atgatatgta	ataatgatat	gtcaaagtat	ttgtattttat	300
ggtcatttaa	ataattaata	atttaattaa	ttttaaatat	tataagaggt	gttaaatatg	360
ccatttggtta	ataaacaatt	taattataaa	gatcctgtaa	atggtgttga	tattgcttat	420
ataaaaaattc	caaatgcagg	acaaatgcaa	ccagtaaaag	cttttaaaat	tcataataaa	480
atatgggtta	ttccagaaag	agatacattt	acaaatcctg	aagaaggaga	tttaaatacca	540
ccaccagaag	caaaacaagt	tccagtttca	tattatgatt	caacatattt	aagtacagat	600
aatgaaaaag	ataattattt	aaaggggagt	acaaaattat	ttgagagaat	ttattcaact	660
gatcttggaa	gaatgttgtt	aacatcaata	gtaaggggaa	taccattttg	gggtggaagt	720
acaatagata	cagaattaaa	agttattgat	actaattgta	ttaatgtgat	acaaccagat	780
ggtagttata	gatacagaaga	acttaatcta	gtaataatag	gaccctcagc	tgataattata	840
cagtttgaat	gtaaaagctt	tggaacatgaa	gttttgaatc	ttacgcgaaa	tggttatggc	900
tctactcaat	acatttagatt	tagcccagat	tttacatttg	gttttgagga	gtcacttgaa	960
gttgatacaa	atcctctttt	aggtgcaggc	aaatttgcta	cagatccagc	agtaacatta	1020
gcacatgaac	ttatacatgc	tggaacataga	ttatatggaa	tagcaattaa	tccaaatagg	1080
gtttttaaag	taaatactaa	tgcctattat	gaaatgagtg	ggttagaagt	aagctttgag	1140
gaaacttagaa	catttggggg	acatgatgca	aagtttatag	atagtttaca	ggaaaacgaa	1200
tttcgtctat	attattataa	taagtttaaa	gatatagcaa	gtacacttaa	taaagctaaa	1260
tcaatagtag	gtactactgc	ttcattacag	tatatgaaaa	atgtttttta	agagaaatat	1320
ctcctatctg	aagatacatc	tggaataattt	tcggtagata	aattaaaaatt	tgataagtta	1380
tacaaaatgt	taacagagat	ttacacagag	gataattttg	ttaagttttt	taaagtactt	1440
aacagaaaaa	catattttgaa	ttttgataaa	gccgtattta	agataaatat	agtacctaag	1500
gtaaattaca	caatatatga	tggaatttaat	ttaagaaata	caaatttagc	agcaaacttt	1560
aatgggtcaaa	atacagaat	taataatatg	aattttacta	aactaaaaaa	ttttactgga	1620
ttgtttgaat	tttataagtt	gctatgtgta	agagggataa	taacttctaa	aactaaatca	1680
ttagataaag	gatacaataa	ggcatttaaat	gattttatgta	tcaaagttta	taattgggac	1740
ttgttttttta	gtccttcaga	agataatttt	actaatgatc	taaataaagg	agaagaaatt	1800
acatctgata	ctaatataga	agcagcagaa	gaaaatatta	gttttagattt	aatacaacaa	1860
tattattttta	ccttttaattt	tgataatgaa	cctgaaaata	tttcaataga	aaatctttca	1920
agtgacatta	tagggccaatt	agaacttatg	cctaatatag	aaagattttcc	taatggaaaa	1980
aagtatgagt	tagataaata	tactatgttc	cattatcttc	gtgctcaaga	atttgaacat	2040
ggtaaatcta	ggattgtctt	aacaaattct	gttaacgaag	cattatttaa	tcctagtcgt	2100
gtttatacat	ttttttcttc	agactatgta	aagaaagtta	ataaagctac	ggaggcagct	2160
atgttttttag	gctgggtaga	acaattagta	tatgatttta	ccgatgaaac	tagcgaagta	2220
agtactacgg	ataaaattgc	ggatataact	ataattattc	catatatagg	acctgcttta	2280
aatataggta	atatgtttata	taaagatgat	tttgtaggtg	ctttaatat	ttcaggagct	2340
gttattctgt	tagaatttat	accagagatt	gcaatacctg	tattaggtac	ttttgcactt	2400
gtatcatata	ttgcgaataa	ggttctaacc	gttcaaacaa	tagataatgc	tttaagtaaa	2460
agaaatgaaa	aatgggatga	ggtctataaa	tatatagtaa	caaattggtt	agcaaagggt	2520
aatacacaga	ttgatctaat	aagaaaaaaa	atgaaagaag	ctttagaaaa	tcaagcagaa	2580
gcaacaaagg	ctataataaa	ctatcagtat	aatcaatata	ctgaggaaga	gaaaaataat	2640
attaatttta	ataattgatga	tttaagttcg	taatttaattg	agtctataaa	taaagctatg	2700
attaatataa	ataaattttt	gaatcaatgc	tctgtttcat	atttaatgaa	ttctatgatc	2760
ccttatgggtg	ttaaacgggt	agaagatttt	gatgctagtc	ttaaagatgc	attattaaag	2820
tatatatatg	ataatagagg	aacttttaatt	ggtcaagtag	atagattaaa	agataaagtt	2880
aataatacac	ttagtacaga	tatacctttt	cagctttcca	aatacgtaga	taatcaaaga	2940
ttattatcta	cattttactga	atatattaag	aatattatta	atacttctat	attgaattta	3000
agatatgaaa	gtaatcattt	aatagactta	tctaggtatg	catcaaaaat	aatattgggt	3060
agtaaagtaa	attttgatcc	aatagataaa	aatcaaatcc	aattatttta	tttagaaagt	3120
agtaaaattg	aggtaatttt	aaaaaatgct	attgtatata	atagtatgta	tgaaaatttt	3180
agtactagct	tttggaataag	aatttcctaag	tatttttaaca	gtataagtc	aaataatgaa	3240

tatacaataa	taaattgtat	ggaaaataat	tcaggatgga	aagtatcact	taattatggt	3300
gaaataatct	ggactttaca	ggatactcag	gaaataaaaac	aaagagtagt	ttttaaatat	3360
agtcaaata	ttaatatatc	agattatata	aacagatgga	tttttgtaac	tatcactaat	3420
aatagattaa	ataactctaa	aatttatata	aatggaagat	taatagatca	aaaaccaatt	3480
tcaaatntag	gtaatatcca	tgctagtaat	aatataatgt	ttaaattaga	tggttgtaga	3540
gatacacata	gatatatttg	gataaaatat	tttaattctt	ttgataagga	attaaatgaa	3600
aaagaaatca	aagattttata	tgataatcaa	tcaaattcag	gtatttttaa	agacttttgg	3660
ggtgattatt	tacaatatga	taaaccatac	tatatgttaa	atttatatga	tccaaataaa	3720
tatgtcgatg	taaataatgt	aggtattaga	ggttatatgt	atcttaaagg	gcctagaggt	3780
agcgtaatga	ctacaaacat	ttattttaat	tcaagtttgt	atagggggac	aaaattttatt	3840
ataaaaaaat	atgcttcttg	aaataaagat	aatattgtta	gaaataatga	tcgtgtatat	3900
attaatgtag	tagttaaaaa	taaagaatat	aggtagctga	ctaatagcac	acaggcaggc	3960
gtagaaaaaa	tactaagtgc	attagaaata	cctgatgtag	gaaatctaag	tcaagtagta	4020
gtaatgaagt	caaaaaatga	tcaaggaata	acaaataaat	gcaaaatgaa	tttacaagat	4080
aataatggga	atgatatagg	ctttatagga	tttcatcagt	ttaataatat	agctaaacta	4140
gtagcaagta	attggtataa	tagacaaata	gaaagatcta	gtaggacttt	gggttgctca	4200
tggaatatta	ttcctgtaga	tgatggatgg	ggagaaaggc	cactgtaatt	aatctcaaac	4260
tacatgagtc	tgtcaagaat	tttctgtaaa	catccataaa	aattttaaaa	ttaatatggt	4320
taagaataac	tagatatgag	tattgtttga	actgcccctg	tcaagtagac	aggtaaaaaa	4380
ataaaaaatta	agatactatg	gtctgatttc	gatattctat	cggagtcaga	ccttttaact	4440
tttcttgtag	cctttttgta	ttgtaaaact	ctatgtattc	atcaattgca	agttccaatt	4500
agtcaaaaatt	atgaaacttt	ctaagataat	acatttctga	ttttataatt	tccaaaaatc	4560
cttccatagg	accattatca	atacatctac	caactcgaga	catactttga	gttgcgctca	4620
tctcattaag	tttattcttg	aaagatttac	ttgtatattg	aaaaccgcta	tcactgtgaa	4680
aaagtggact	agcatcagga	ttggaggtaa	ctgctttatc	aaaggtttca	aagacaagga	4740
cgttgttatt	tgattttcca	agtacatagg	aaataatgct	attatcatgc	aatcaagta	4800
tttcaactcaa	gtacgccttt	gtttcgtctg	ttaac			4835

<210> 11

<211> 15

<212> PRT

<213> Homo sapiens

<400> 11

Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro
1				5					10				15	